

Supplemental Table S1. Changes in Age at Death among Control and Diabetic Individuals with Comorbidities

Year	Control		Diabetes (n=601,990)							
			Group I: HTN(-), DL(-)		Group II: HTN(-), DL(+)		Group III: HTN(+), DL(-)		Group IV: HTN(+), DL(+)	
	No.	Mean±SD	No.	Mean±SD	No.	Mean±SD	No.	Mean±SD	No.	Mean±SD
Overall	830,577		97,522		28,089		287,605		188,774	
2006	79,224	76.0±14.3	7,077	67.9±11.9	568	66.6±11.1	17,331	72.2±9.82	4,666	70.5±9.2
2007	66,965	76.3±14.6	7,402	67.8±12.1	736	66.4±11.3	20,123	72.6±10.2	6,443	70.9±9.5
2008	62,301	76.4±14.8	7,184	67.9±12.3	897	67.3±11.3	21,259	73.0±10.2	7,736	71.3±9.6
2009	58,532	76.2±15.0	7,183	68.4±12.3	1,064	66.9±11.8	21,448	73.7±10.2	8,864	71.8±9.7
2010	60,658	76.6±14.9	7,313	68.7±12.3	1,310	67.4±11.9	22,233	74.1±10.3	10,415	72.4±9.7
2011	59,873	76.9±14.8	7,207	68.9±12.2	1,484	67.6±11.5	22,815	74.6±10.4	11,921	72.9±9.7
2012	62,098	77.2±14.7	7,629	69.7±12.3	1,784	69.3±11.3	23,752	75.3±10.3	14,070	73.7±9.7
2013	59,356	77.2±14.7	7,557	70.1±12.4	2,131	69.4±11.8	23,844	75.6±10.3	15,412	74.1±9.7
2014	59,972	77.5±14.6	7,649	70.4±12.3	2,411	70.0±11.4	23,473	76.0±10.4	17,172	74.5±9.7
2015	62,425	78.0±14.5	7,739	71.0±12.2	2,981	70.5±11.6	23,025	76.7±10.4	19,564	75.1±9.7
2016	63,661	78.2±14.3	7,703	71.3±12.4	3,647	71.1±11.5	22,632	77.0±10.4	21,812	75.5±9.7
2017	66,191	78.8±14.2	7,826	71.8±12.4	4,120	72.0±11.4	22,480	77.4±10.5	23,825	76.1±9.8
2018	69,321	79.1±14.0	8,053	72.5±12.3	4,956	72.6±11.4	23,190	77.8±10.5	26,874	76.4±9.8
Men	415,166		66,610		17,281		157,240		98,353	
2006	39,330	70.4±14.5	4,688	65.3±11.5	309	63.8±11.3	9,263	69.8±10.0	2,232	68.6±9.4
2007	33,933	70.7±14.8	4,881	65.1±11.5	430	63.4±11.0	10,637	69.8±10.5	3,040	68.6±9.7
2008	32,163	70.7±14.9	4,907	65.5±11.8	549	64.9±11.1	11,609	70.4±10.5	3,741	69.0±9.8
2009	30,444	70.5±15.0	4,878	65.6±11.8	613	64.0±11.1	11,669	70.9±10.5	4,404	69.4±9.9
2010	31,048	70.9±15.0	5,047	66.3±11.7	775	64.8±11.8	12,236	71.3±10.6	5,192	70.0±10.0
2011	30,407	71.1±14.9	4,951	66.6±11.8	923	65.4±11.2	12,587	71.8±10.6	6,122	70.7±9.9
2012	31,330	71.4±14.9	5,287	67.3±11.7	1,081	67.3±11.2	13,062	72.3±10.4	7,134	71.4±9.9
2013	29,824	71.4±14.9	5,212	67.9±11.9	1,278	67.0±11.5	13,096	72.7±10.5	7,988	71.8±10.0
2014	30,041	71.7±14.7	5,278	68.0±11.8	1,490	68.1±11.2	12,955	72.9±10.6	9,122	72.4±9.9
2015	30,956	72.1±14.7	5,371	68.9±11.8	1,811	68.6±11.3	12,528	73.6±10.6	10,414	72.9±10.0
2016	31,124	72.5±14.5	5,299	69.0±12.0	2,299	69.5±11.4	12,379	74.0±10.5	11,599	73.4±9.8
2017	31,633	72.7±14.5	5,309	69.6±11.9	2,624	70.4±11.0	12,380	74.4±10.5	12,783	73.8±10.0
2018	32,933	73.3±14.4	5,502	70.4±11.7	3,099	71.1±11.1	12,839	74.8±10.6	14,582	74.1±9.9
Women	415,411		30,912		10,808		130,365		90,421	
2006	39,894	81.5±11.6	2,389	72.8±10.9	259	70.0±9.8	8,068	75.0±8.8	2,434	72.3±8.6
2007	33,032	82.1±11.9	2,521	72.9±11.6	306	70.6±10.4	9,486	75.7±8.9	3,403	72.3±8.8
2008	30,138	82.4±12.1	2,277	72.9±11.8	348	71.2±10.6	9,650	76.2±8.9	3,995	72.3±9.0
2009	28,088	82.4±12.3	2,305	74.2±11.2	451	70.8±11.6	9,779	77.0±8.7	4,460	72.3±8.9
2010	29,610	82.5±12.2	2,266	74.1±11.7	535	71.3±10.9	9,997	77.6±8.8	5,223	72.3±8.7
2011	29,466	82.8±12.0	2,256	74.0±11.7	561	71.3±11.0	10,228	78.1±9.0	5,799	72.3±8.9
2012	30,768	83.2±11.9	2,342	75.0±11.8	703	72.4±10.8	10,690	78.9±8.8	6,936	72.3±8.9
2013	29,532	83.0±12.0	2,345	75.0±12.1	853	73.0±11.4	10,748	79.2±9.0	7,424	72.3±8.7
2014	29,931	83.3±12.0	2,371	75.6±11.8	921	73.1±11.0	10,518	79.7±8.8	8,050	72.3±9.0
2015	31,469	83.7±11.8	2,368	75.7±11.8	1,170	73.4±11.4	10,497	80.3±8.9	9,150	72.3±8.8
2016	32,537	83.7±11.7	2,404	76.3±11.8	1,348	73.9±11.1	10,253	80.6±9.0	10,213	72.3±8.9
2017	34,558	84.3±11.3	2,517	76.3±12.2	1,496	74.7±11.5	10,100	81.1±9.1	11,042	72.3±8.8
2018	36,388	84.4±11.4	2,551	77.1±12.1	1,857	75.2±11.4	10,351	81.5±9.1	12,292	72.3±8.9

HTN, hypertension; DL, dyslipidemia; SD, standard deviation.