EnM

Study name	Statistics with study removed			Risk ratio (95%) by
	Risk ratio	Lower limit	Upper limit	study removed
Ahn et al. 2011	0.247	0.126	0.483	-■-
Kashima et al. 1998	0.307	0.147	0.642	
Kwak et al. 2014	0.233	0.119	0.455	
Kebebew et al. 2001	0.254	0.128	0.505	
NEST (Present study)	0.159	0.058	0.441	
				0.1 1 10 A

Study name	Statistics with study removed			Risk ratio (95%) by
	Risk ratio	Lower limit	Upper limit	study removed
Xu et al. 2021	0.259	0.143	0.467	-■-
Huang et al. 2011	0.200	0.111	0.361	-■-
Ahn et al. 2011	0.221	0.127	0.385	-■-
Loh et al. 1999	0.248	0.141	0.436	
Kashima et al. 1998	0.256	0.142	0.462	-■-
Kebebew et al. 2001	0.239	0.137	0.416	-■-
NEST (Present study)	0.179	0.088	0.363	 ■

Supplemental Fig. S3. Sensitivity analysis between Hashimoto thyroiditis and mortality. (A) All-cause mortality. (B) Thyroid cancer-related mortality. NEST, National Epidemiologic Survey of Thyroid cancer.